

Table 3G
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area

Page: 1
Run Date: 7 / 12 / 11

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dane Cty	2,052	39.9	23.5	25.9	0.4	7.5	2.8	0.0
Mil. Cty	9,862	17.1	49.5	26.6	0.5	4.5	1.8	0.0
Racine	1,400	41.1	25.2	30.5	0.1	0.6	2.4	0.0
Rest of WI	20,475	73.1	3.5	13.6	2.7	4.8	2.3	0.0
Wisconsin	33,789	53.4	19.0	18.9	1.8	4.7	2.2	0.0
Nation (Prior Year)	1,304,320	41.6	24.1	28.3	1.1	3.2	1.2	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4G
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Metro Area

Comparison of Age Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	2,052	0.3	4.5	9.9	59.4	24.3	1.4	0.1
Mil. Cty	9,862	0.4	6.8	12.6	58.8	20.1	1.3	0.1
Racine	1,400	0.6	6.6	12.6	60.1	19.0	1.2	0.0
Rest of WI	20,475	0.1	4.2	12.6	63.1	18.6	1.3	0.0
Wisconsin	33,789	0.2	5.1	12.5	61.5	19.4	1.3	0.0
Nation (Prior Year)	1,304,320	0.3	5.8	12.3	59.4	20.4	1.5	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5G
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Metro Area

Page: 3
Run Date: 7 / 12 / 11

Comparison of Maternal Health Indicators ^(2, 3)															
Metro Area	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Dane Cty	2,022	2.9 (1)	56.8 (2)	1,487	22.0 (3)	50.8 (3)	345	29.6 (2)	1,655	21.8 (2)	1,833	6.2 (4)	1,833	5.0(1)	
Mil. Cty	9,761	3.2 (2)	59.9 (4)	8,255	25.6 (4)	46.6 (1)	1,576	44.7 (4)	8,288	32.5 (4)	9,040	4.9 (1)	9,040	6.6(3)	
Racine	1,386	3.4 (3)	57.9 (3)	1,247	21.0 (1)	51.8 (4)	205	33.2 (3)	1,128	25.4 (3)	1,273	5.5 (3)	1,273	7.4(4)	
Rest of WI	20,246	3.6 (4)	55.0 (1)	15,667	21.6 (2)	50.2 (2)	3,183	26.1 (1)	16,027	19.6 (1)	18,597	5.2 (2)	18,597	5.4(2)	
Wisconsin	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9	30,743	5.2	30,743	5.8	
Nation (Prior Year)	1,226,875	4.5	52.9	917,032	21.2	48.2	189,944	33.9	412,633	28.8	389,161	5.0	394,368	6.8	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6G
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Metro Area	Medical Care					WIC Enrollment			Multivitamin Consumption	
	1st Trimester		No Care			1st Trimester			Prepregnancy (4)	During Pregnancy (5)
	Number	% (Rank)	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Dane Cty	1,847	91.9 (1)	1.2 (1)			2,044	33.4 (4)		1,832	19.7 (2)
Mil. Cty	9,222	83.5 (4)	2.7 (3)			9,827	35.2 (3)		9,666	12.2 (4)
Racine	1,266	83.6 (3)	2.9 (4)			1,397	39.2 (1)		1,377	17.9 (3)
Rest of WI	18,893	88.7 (2)	2.5 (2)			20,403	36.2 (2)		19,815	21.5 (1)
Wisconsin	31,228	87.2	2.5			33,671	35.8		32,690	18.5
Nation (Prior Year)	527,184	82.2	3.1			1,213,982	34.1		360,369	18.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7G
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Metro Area

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	1,983	28.6 (2)	1,833	15.2 (2)	415	39.3 (1)	1,638	12.8 (1)	1,833	3.6 (1)
Mil. Cty	9,161	22.9 (1)	9,036	11.5 (1)	1,637	38.9 (2)	7,856	18.4 (3)	9,040	6.8 (3)
Racine	1,350	31.6 (3)	1,273	15.6 (3)	328	35.4 (3)	1,132	21.3 (4)	1,273	9.0 (4)
Rest of WI	19,554	40.7 (4)	18,565	22.3 (4)	6,088	30.5 (4)	16,305	17.2 (2)	18,574	4.5 (2)
Wisconsin	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3
Nation (Prior Year)	1,040,568	22.9	838,781	13.0	156,709	36.7	650,648	17.1	559,995	16.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8G
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Metro Area

Comparison of Infant Health Indicators (2, 3)										
Metro Area	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	1,779	7.9 (3)	8.3 (4)		1,846	9.3 (1)	1,616	3.2 (3)	1,810	81.9 (1)
Mil. Cty	8,728	9.7 (4)	5.6 (1)		9,050	11.0 (4)	7,772	4.2 (4)	8,838	65.3 (3)
Racine	1,215	7.2 (2)	6.3 (2)		1,268	9.7 (3)	1,097	2.4 (1)	1,236	64.6 (4)
Rest of WI	17,962	7.1 (1)	7.7 (3)		18,620	9.4 (2)	16,247	2.6 (2)	18,247	72.2 (2)
Wisconsin	29,684	7.9	7.1		30,784	9.9	26,732	3.1	30,131	70.4
Nation (Prior Year)	1,018,145	8.2	6.7		1,017,333	11.1	848,957	3.5	997,136	66.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3G
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dane Cty	6,370	41.2	22.1	26.2	0.5	7.5	2.5	0.0
Mil. Cty	31,067	17.7	49.6	26.2	0.6	4.4	1.6	0.0
Racine	4,475	41.2	26.7	29.1	0.2	0.6	2.3	0.0
Rest of WI	63,322	73.3	3.3	14.0	2.7	4.6	2.1	0.0
Wisconsin	105,234	53.6	19.1	19.0	1.8	4.6	2.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4G
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Metro Area
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	6,370	0.3	4.7	10.7	60.1	22.7	1.4	0.0
Mil. Cty	31,067	0.5	7.6	13.2	58.8	18.5	1.3	0.0
Racine	4,475	0.6	6.8	12.8	61.5	17.2	1.1	0.0
Rest of WI	63,322	0.2	4.6	13.1	63.2	17.8	1.2	0.0
Wisconsin	105,234	0.3	5.6	13.0	61.6	18.3	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5G
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Metro Area
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)															
Metro Area	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Dane Cty	6,276	3.2 (1)	53.5 (1)	4,199	21.4 (2)	50.2 (2)	1,180	28.1 (2)	4,822	20.2 (1)	5,684	5.9 (4)	5,684	5.3(2)	
Mil. Cty	30,741	3.2 (1)	57.8 (4)	23,020	23.8 (4)	48.8 (1)	5,050	44.0 (4)	25,998	33.8 (4)	28,342	4.4 (1)	28,342	6.7(3)	
Racine	4,430	3.2 (1)	57.5 (3)	3,959	21.4 (2)	52.0 (4)	733	34.2 (3)	3,667	24.7 (3)	4,056	4.6 (3)	4,056	6.7(3)	
Rest of WI	62,588	3.9 (4)	53.9 (2)	46,103	20.7 (1)	51.3 (3)	10,026	26.6 (1)	50,437	20.2 (1)	57,553	4.5 (2)	57,553	5.2(1)	
Wisconsin	104,035	3.6	55.2	77,281	21.7	50.5	16,989	32.2	84,924	24.6	95,635	4.6	95,635	5.7	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6G
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Metro Area	Medical Care					WIC Enrollment			Multivitamin Consumption	
	1st Trimester		No Care			1st Trimester			Prepregnancy (4)	During Pregnancy (5)
	Number	% (Rank)	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Dane Cty	5,730	88.8 (1)	1.3 (1)			6,281	29.2 (4)		5,839	17.1 (2)
Mil. Cty	28,903	81.1 (4)	2.8 (4)			30,611	34.1 (3)		30,441	12.0 (4)
Racine	4,056	82.5 (3)	2.7 (3)			4,429	37.0 (1)		4,386	15.1 (3)
Rest of WI	58,186	87.7 (2)	2.5 (2)			62,561	34.7 (2)		61,166	20.1 (1)
Wisconsin	96,875	85.6	2.5			103,882	34.3		101,832	17.3
									82,891	83.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7G
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Metro Area
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	6,134	29.5 (2)	5,674	15.6 (2)	1,263	38.5 (1)	5,038	14.1 (1)	5,675	4.3 (1)
Mil. Cty	29,115	23.8 (1)	28,332	12.2 (1)	5,247	37.5 (2)	24,887	20.8 (3)	28,335	8.9 (4)
Racine	4,295	32.9 (3)	4,054	16.3 (3)	1,080	33.0 (3)	3,654	22.3 (4)	4,054	8.7 (3)
Rest of WI	60,569	42.3 (4)	57,430	23.8 (4)	19,688	29.6 (4)	50,380	19.8 (2)	57,470	5.7 (2)
Wisconsin	100,113	35.7	95,490	19.5	27,278	31.7	83,959	19.8	95,534	6.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8G
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Metro Area
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
Metro Area	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	5,483	6.9 (1)	8.9 (4)		5,654	9.1 (1)		4,928	2.7 (2)	5,583 80.5 (1)
Mil. Cty	27,288	9.5 (4)	6.0 (1)		28,027	10.9 (4)		24,018	4.0 (4)	27,655 65.0 (3)
Racine	3,910	8.8 (3)	6.3 (2)		4,016	10.7 (3)		3,458	3.2 (3)	3,955 62.0 (4)
Rest of WI	55,575	7.0 (2)	8.3 (3)		57,091	9.4 (2)		49,839	2.6 (1)	56,446 69.9 (2)
Wisconsin	92,256	7.8	7.6		94,788	9.9		82,243	3.1	93,639 68.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

